arXiv:hep-ph/9903345v1 15 Mar 1999

The rare decays $B \to X_{s,d}\ \nu\bar{\nu}$ and $B_{s,d} \to l^+l^-$ in the Multiscale Walking Technicolor Model *

Xiao Zhenjun[1,2], Lü Linxia[1], Guo Hongkai[1], Lu Gongru[1]

[1] Department of Physics, Henan Normal University, Xinxiang, 453002

[2] Department of Physics, Peking University, Beijing, 100871

February 1, 2008

Abstract

We calculate the contributions to the rare B-decays, $B \to X_{s,d}\nu\bar{\nu}$, $B_{s,d} \to l^+l^-$ from the unit-charged technipions. Within the considered parameter space we find that: (a) the enhancements to the branching ratios in question can be as large as three orders of magnitude; (b) the ALEPH data of $B \to X_s\nu\bar{\nu}$ lead to strong mass bounds on m_{p1} and m_{p8}: $m_{p8} \geq 620, 475GeV$ for $F_Q = 40GeV$ and $m_{p1} = 100, 400GeV$ respectively. (c) the CDF data of $B_s \to \mu\bar{\mu}$ lead to a relatively weak limit: $m_{p8} \geq 320GeV$ for $F_Q = 40GeV$ and $m_{p1} = 200GeV$.

PACS: 12.60.Nz, 12.15.Ji, 13.20.Jf

In the framework of the Standard Model (SM), the rare decays $B \to X_{s,d}\ \nu\overline{\nu}$ and $B_{s,d} \to l^+l^-$ are theoretically very clean and dominated by similar Z^0-penguin and W-box diagrams involving top quark exchanges [1, 2]. These rare B-decay modes therefore may play an important role in searching for the new physics beyond the SM [3].

In ref.[4], Lane and Ramana constructed a specific multiscale walking technicolor model(MWTCM) and investigated its phenomenology. This model predicted a rich spectrum of technipions. Among them are unit-charged color-octets $\pi^a_{\bar{D}U}$ (a is the color index) and unit-charged color-singlets P_1^+ and P_2^+. In this letter, we use the symbol π_8 and m_{p8} to denote the color-octet $\pi_{\bar{D}U}$ and its mass. The mixed state π_1 of the P_1^+ and P_2^+ is the same kind of technipion as P^+ given in ref.[5]. We will study the new contributions to

*Supported by the National Natural Science Foundation of China under Grant No.19575015 and By the Sino-British Friendship Scholarship Scheme.

the rare B-decays from the physical mixed state π_1 instead of the two technipions P_1^+ and P_2^+, for the sake of simplicity. According to the estimations in ref.[4], $m_{p1} \sim 200GeV$ and $m_{p8} \sim 300GeV$.

In ref.[6], the authors calculated the contributions to the rare decay $b \to s\gamma$ due to the effective $bs\gamma$ coupling induced by the π_1 and π_8 appeared in the MWTCM and found that resultant enhancement to the branching ratio $B(B \to X_s\gamma)$ can be two orders of magnitude. The CLEO data[7] led to strong limit on charged technipion masses: $m_{p8} > 600GeV$ for $m_{p1} = 300GeV$. The MWTCM itself is therefore strongly disfavored by the CLEO data[7].

In this letter, we calculate the new contributions to the rare decays $B \to X_{s,d}\,\nu\overline{\nu}$ and $B_{s,d} \to l^+l^-$ due to the effective $b\overline{s}Z$ coupling induced also by the π_1 and π_8. This work is complementary to the relevant studies about the new effects to the $b \to s\gamma$ decay[6] and the rare K-decays *etc* [8, 3], in order to test or constrain the Technicolor(TC) models by currently available data of rare B- and K-decays.

In the numerical calculation, we treat m_{p1} and m_{p8} as semi-free parameters, varying in the ranges of $100GeV \leq m_{p1} \leq 400GeV$ and $200GeV \leq m_{p8} \leq 800GeV$ respectively. The relevant Yukawa and gauge couplings of charged technipions to fermion pairs and to Z^0 gauge boson can be found in refs.[3, 5, 9].

The new penguin diagrams for the induced $b\overline{s}Z$ couplings due to the exchange of the π_1 and π_8 are shown in Fig.1. The corresponding one-loop diagrams in the SM were evaluated long time ago and can be found in ref.[10]. Only the color-singlet π_1 couples to lepton pairs, and therefore may contribute to the rare B-decays in question through the box diagrams. But the Yukawa couplings between π_1 and $l\nu$ pairs are strongly suppressed by the lightness of the lepton masses m_l ($l = e\mu, \tau$). Consequently, we can neglect the tiny contributions from π_1 through the box diagrams safely.

Because of the lightness of the s and b quarks when compared with the top quark mass m_t and the technipion masses m_{p1} and m_{p8} we set $m_s = 0$ and $m_b = 0$ in the calculation. We will use dimensional regularization to regulate all the ultraviolet divergences in the virtual loop corrections and adopt the modified minimal subtracted ($\overline{MS}$) renormalization scheme.

By analytical evaluations of the Feynman diagrams as shown in Fig.1, we find the effective $b\overline{s}Z$ vertex induced by the π_1 and π_8 exchanges,

$$\Gamma^I_{Z_\mu} = \frac{1}{16\pi^2}\frac{g^3}{\cos\theta_W}\ \sum_j V^*_{js}V_{jb}\,\overline{s_L}\,\gamma_\mu\,b_L\ C_0^{New}(y_j) \tag{1}$$

$$\Gamma^{II}_{Z_\mu} = \frac{1}{16\pi^2}\frac{g^3}{\cos\theta_W}\ \sum_j V^*_{js}V_{jb}\,\overline{s_L}\,\gamma_\mu\,b_L\ C_0^{New}(z_j) \tag{2}$$

with

$$C_0^{New}(y_j) = \eta^a_{TC}\ \left[\frac{y_j(-1+2\sin^2\theta_W-3y_j+2\sin^2\theta_W y_j)}{8(1-y_j)} - \frac{\cos^2\theta_W y_j^2}{2(1-y_j)^2}\ln(y_j)\right] \tag{3}$$

$$C_0^{New}(z_j)=\eta_{TC}^b\left[\frac{z_j(-1+2\sin^2\theta_W-3z_j+2\sin^2\theta_W z_j)}{8(1-z_j)}-\frac{\cos^2\theta_W z_j^2}{2(1-z_j)^2}\ln(z_j)\right] \quad (4)$$

and

$$\eta_{TC}^a=\frac{m_{p1}^2}{3\sqrt{2}F_Q^2G_FM_W^2},\quad \eta_{TC}^b=\frac{8m_{p8}^2}{3\sqrt{2}F_Q^2G_FM_W^2} \quad (5)$$

where $y_j=m_j^2/m_{p1}^2$, $z_j=m_j^2/m_{p8}^2$, V_{ij} ($i=u,c,t$ and $j=d,s,b$) are the elements of the CKM mixing matrix, M_W is the mass of W gauge boson, F_Q is the technipion decay constant in the MWTCM, $\sin\theta_W$ is the Weinberg angle, and $G_F=1.16639\times 10^{-5}(GeV^{-2})$ is the Fermi coupling constant. The functions $C_0^{New}(y_j)$ and $C_0^{New}(z_j)$ in eqs.(3, 4) are just the same kind of functions as the basic function $C_0(x_i)$ in eq.(2.18) of ref.[1]. Functions $C_0(y_t)$ and $C_0(z_t)$ describe the contributions to the $b\overline{s}Z$ vertex from the π_1 and π_8 respectively. The $C_0(y_t)$ is always positive, but the $C_0(z_t)$ will change its sign from " $+$ " to " $-$ " at $m_{p8}=531GeV$.

Within the SM, the rare B-decays under consideration depend on the functions $X(x_t)$ and/or $Y(x_t)$, they are currently known at the next-to-leading order level [1, 2]. When the new contributions from charged technipions are included, one has

$$X(x_t,y_t,z_t) = X(x_t)+C_0^{New}(y_t)+C_0^{New}(z_t), \quad (6)$$
$$Y(x_t,y_t,z_t) = Y(x_t)+C_0^{New}(y_t)+C_0^{New}(z_t). \quad (7)$$

where $x_t=m_t^2/m_W^2$, $y_t=m_t^2/m_{p1}^2$ and $z_t=m_t^2/m_{p8}^2$.

In the following numerical calculations, we fix the relevant parameters as follows and use them as the Standard Input: $M_W=80.2GeV$, $\alpha_{em}=1/129$, $\sin^2\theta_W=0.23$, $m_t\equiv\overline{m_t}(m_t)=170GeV$, $\tau(B_s)=\tau(B_d)=1.6ps$, $\Lambda_{\overline{MS}}^{(5)}=0.225GeV$, $F_{B_s}=0.210GeV$, $m_{B_s}=5.38GeV$, $m_{B_d}=5.28GeV$, $A=0.84$, $\lambda=0.22$, $\rho=0$ and $\eta=0.36$. For the definitions and values of these input parameters, one can see refs.[1, 2].

Within the SM, normalizing to the semi-leptonic branching ratio $B(B\to X_ce\overline{\nu})$ and summing over the three neutrino flavors one finds[1]

$$B(B\to X_s\,\nu\overline{\nu})=B(B\to X_ce\overline{\nu})\frac{3\alpha_{em}^2}{4\pi^2\sin^4\theta_W}\frac{\mid V_{ts}\mid^2}{\mid V_{cb}\mid^2}\frac{X(x_t)^2}{f(z)}\frac{\overline{\eta}}{\kappa(z)} \quad (8)$$

where $\bar{\eta}=\kappa(0)$, $f(z)$ and $\kappa(z)$ with $z=m_c/m_b=0.29$ are the phase-space and quantum chromodynamics(QCD) correction factors for the decay $B\to X_ce\overline{\nu}$,

$$f(z) = 1-8z^2+8z^6-z^8-24z^4\ln(z), \quad (9)$$
$$\kappa(z) = 1-\frac{2\alpha_s(m_b)}{3\pi}\left[(\pi^2-\frac{31}{4})(1-z)^2+\frac{3}{2}\right] \quad (10)$$

where $\alpha_s(m_b)$ is the QCD coupling constant at the energy scale $\mu=m_b$.

Table 1: The branching ratios of $B_{s,d} \to l^+l^-$ for l=e,μ,τ.

Branching ratio	SM	SM + New	Data
$B(B_s \to e^+e^-)$	0.73×10^{-13}	$1.9 \times 10^{-10} - 1.7 \times 10^{-12}$	
$B(B_s \to \mu^+\mu^-)$	3.13×10^{-9}	$8.1 \times 10^{-6} - 7.3 \times 10^{-8}$	$\leq 2.0 \times 10^{-6}$
$B(B_s \to \tau^+\tau^-)$	0.67×10^{-6}	$1.7 \times 10^{-3} - 1.5 \times 10^{-5}$	
$B(B_d \to e^+e^-)$	0.44×10^{-14}	$1.2 \times 10^{-12} - 1.0 \times 10^{-14}$	
$B(B_d \to \mu^+\mu^-)$	1.90×10^{-10}	$4.9 \times 10^{-8} - 4.4 \times 10^{-10}$	$\leq 6.8 \times 10^{-7}$
$B(B_d \to \tau^+\tau^-)$	0.40×10^{-7}	$1.1 \times 10^{-5} - 0.9 \times 10^{-7}$	

Within the SM, using the Standard input parameters and setting $B(B \to X_c e\overline{\nu}) = 10.4\%$ and $\mid V_{ts}/V_{cb} \mid^2 = 0.95$, one finds $B(B \to X_s\nu\bar{\nu}) = 3.52 \times 10^{-5}$ and $B(B \to X_d\nu\bar{\nu}) = 2.03 \times 10^{-6}$. When the new contributions due to π_1 and π_8 are included, one has $B(B \to X_s\nu\bar{\nu}) = 1.57 \times 10^{-3}$ for the typical values of $F_Q = 40GeV$, $m_{p1} = 200GeV$ and $m_{p8} = 300GeV$, that is two-orders of magnitude higher than the value of the SM prediction. The color-octet π_8 dominates the total contribution.

In Fig.2 the dot-dash (solid, dots) curve shows the branching ratio when the new contributions from π_1 and π_8 are taken into account for $F_Q = 40GeV$ and $m_{p1} = 100GeV$ ($200GeV$, $400GeV$). The horizontal short-dash line shows the ALEPH bound on the branching ratio $B(B \to X_s\nu\bar{\nu})$ [11]: $B(B \to X_s\nu\bar{\nu}) < 7.7 \times 10^{-4}$, which is a factor of 20 above the SM expectation but sensitive enough to put stringent limits on technipion masses. Assuming $F_Q = 40GeV$, one has $m_{p8} > 620GeV$ for $m_{p1} = 100GeV$, $530GeV < m_{p8} < 890GeV$ for $m_{p1} = 200GeV$ and $475GeV < m_{p8} < 740GeV$ for $m_{p1} = 400GeV$. For smaller F_Q, the constraints will become much stronger.

In the case of the decay $B \to X_d\ \nu\overline{\nu}$ one has to replace V_{ts} in eq.(8) by V_{td}, which results in a decrease of the branching ratio by roughly an order of magnitude. But unfortunately, no experimental bound on the decay $B \to X_d\ \nu\overline{\nu}$ is available currently.

Within the SM, using the effective Hamiltonian as given in ref.[1] one finds

$$B(B_s \to l^+l^-) = \frac{\tau(B_s)G_F^2}{\pi}(\frac{\alpha_{em}}{4\pi\sin^2\theta_W})^2 F_{B_s}^2 m_l^2 m_{B_s}\sqrt{1-\frac{4m_l^2}{m_{B_s}^2}} \mid V_{tb}^*V_{ts} \mid^2 Y(x_t)^2 \qquad (11)$$

with s replaced by d in the case of $B_d \to l^+l^-$.

In the numerical calculations, we use the standard input parameters and assume that $F_Q = 40GeV$, $m_{p1} = 200GeV$, $200GeV \leq m_{p8} \leq 800GeV$, and set $\mid V_{tb}^*V_{ts} \mid^2 = 0.0021$ and $\mid V_{tb}^*V_{td} \mid^2 = 1.3 \times 10^{-4}$. The numerical results with the inclusion of new contributions from the technipions π_1 and π_8 are listed in Table 1.

For the decays $B_s \to l^+l^-$, the available experimental bound is $B(B_s \to \mu^+\mu^-) \leq 2.0 \times 10^{-6}$[12], which leads to the lower bounds on m_{p8}: $m_{p8} > 320GeV$ for $F_Q = 40GeV$

and $m_{p1} = 200GeV$. But this bound is much weaker than that from the ALEPH data of $B \to X_s \nu\bar{\nu}$. For the decays $B_d \to l^+ l^-$, the available experimental bound is $B(B_d \to \mu^+\mu^-) \leq 6.8 \times 10^{-7}$ [12], which is still not sensitive enough to put any limits on m_{p1} and m_{p8}.

In summary, the ALEPH data of $B \to X_s \nu\bar{\nu}$ lead to strong limits on the charged technipion masses m_{p1} and m_{p8}. The assumed mass ranges of π_1 and π_8 in the MWTCM [4] are excluded and therefore the model itself is strongly disfavored by ALEPH data[11]. Other relevant studies[6, 8] also led to the similar conclusions. The major problem of the MWTCM is that the heavy top quark mass is assumed to be generated by extended technicolor interaction, which is clearly unreasonable[13].

References

[1] A. J. Buras and R.Fleischer, in *Heavy Flavor II*, edited by A.J.Buras and M.Lindner (World Scientific Publishing Co., Singapore, 1998).

[2] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev.Mod.Phys. 68(1996)1125.

[3] C.D. Lü and Z.J. Xiao, Phys.Rev. D53(1996)2529; Yue Chong-xing and Lu Gong-ru, Chin.Phys.Lett. 16(9)(1998)631; Xiao Zhen-jun et al., Commun.Theor.Phys. 29 (1998)431; Xiao Zhen-jun, Zhang Jin-yu, Lü Lin-xia and Lu Gong-ru, *High Energy Phys. and Nucl. Phys. (in Chinese)*, 23(1999) (in press); G.R. Lu, Z.H. Xiong and Y.G. Cao, Nucl.Phys. B487(1997)43.

[4] K. Lane and M.V. Ramana, Phys.Rev. D44(1991)2678.

[5] E. Farhi and L. Susskind, Phys. Rev. D20(1979)3404. E. Eichten, I. Hinchliffe, K. Lane and C. Quigg, Rev.Mod.Phys. 56(1984)579; Phys.Rev. D34(1986)1547.

[6] G.R. Lu, Y.G. Cao, Z.H. Xiong and C.Y. Yue, Z.Phys. C74(1997)355.

[7] M.S. Alam et al., (CLEO Collaboration), Phys. Rev. Lett. 74(1995)2885.

[8] Z.J. Xiao, L.X. Lü, H.K. Guo and G.R. Lu, Eur. Phys. J. C7(1999)487.

[9] J. Ellis et al., Nucl.Phys. B182(1981)505.

[10] T. Inami and C.S. Lim, Prog.Theor.Phys. 65(1981)297.

[11] H.Kroha et al., (ALEPH Collaboration), Proceedings of the 28th International Conference on High Energy Physics, edited by Z.Ajduk and A.K.Wroblewski, (World Scientific Publishing Co., Singapore, 1997), p.1182.

[12] F. Abe et al., (CDF Collaboration), Phys.Rev. D57(1998)R3811.

[13] K.Lane, Proceedings of the 28th International Conference on High Energy Physics, edited by Z.Ajduk and A.K.Wroblewski, (World Scientific Publishing Co., Singapore, 1997), p.367.

Figure Captions

Fig.1: New Z^0-penguin diagrams contributing to the induced $b\bar{s}Z$ vertex from the internal exchanges of the technipion π_1 and π_8. The dashed lines are π_1 and π_8 lines and u_j stands for the quarks (u, c, t).

Fig.2: The m_{p8} dependence of the branching ratio $B(B \to X_s \nu\bar{\nu})$ when the new contributions are included. The horizontal short-dash line shows the ALEPH upper bound, while the dot-dash (solid, dots) curve shows the branching ratio for $m_{p1} = 100GeV$ $(200GeV,\ 400GeV)$.





Fig.1


Br(B -> Xs ν ν̄) (10-5)
200
150
100
50
0
200
400
600
800
MP8(GeV)
FQ = 40GeV
ALEPH Data